


**Provided for non-commercial research and education use.  
Not for reproduction, distribution or commercial use.**




Egyptian Academic Journal of Biological Sciences is the official English language journal of the Egyptian Society of Biological Sciences, Department of Entomology ,Faculty of Sciences Ain Shams University .

The Journal publishes original research papers and reviews from any zoological discipline or from directly allied fields in ecology, behavioral biology, physiology, biochemistry, development, genetics, systematics, morphology, evolution, and general zoology

[www.eajbs.eg.net](http://www.eajbs.eg.net)

---

**Citation:** *Egypt. Acad. J. Biolog. Sci. (B. Zoology) Vol. 7(1)pp 45-52(2015)*



---

## Heavy Metals (Pb, Fe and Zn) in Fish Due to Water Toxicity

El- Sayed A. M. Shokr

Physiology Department, Faculty of Medicine, Hail University, KSA

---

### ARTICLE INFO

#### Article History

Received: 25/3/2015

Accepted: 5/5/2015

---

#### Keywords:

*Oreochromis niloticus*

Pollution

Mmetals

Tissues

### ABSTRACT

Water Toxicity is a worldwide problem, heavy metals belonging to the most important pollutants. The progress of industries has led to increased emission of pollutants into ecosystems. The fish were divided into four groups, control group and three groups exposed to 5, 10 and 15 mg/l of lead acetate, 10, 20 and 30 mg L<sup>-1</sup> of ferric chloride and 10, 20 and 300 mg L<sup>-1</sup> of zinc sulfate respectively, the experimental fish were exposed to treatment for periods of 15 days, 30 days and 45 days. The Nile tilapia was exposed to water pollution with lead, iron and zinc. The Nile tilapia 80 - 100 gm body weight was obtained from Abbassa farm. The fish were reared in large tanks contained water from their farm. The fish were acclimatized and fed on 30 % protein diet. Result of the present study revealed that lead concentrations in the tissues of the fish were elevated and highest contamination were increased in the gill and liver than that in the muscles. Also, there was vary significant increase in the concentrations of lead in the gill, liver and muscles tissues of *Oreochromis niloticus* exposed to different concentrations of lead acetate than that control fish tissues. Iron concentrations in the tissues of the fish were elevated in the gill and liver than that in the muscles. Also, there was vary significant increase in the concentrations of iron in the gill, liver and muscle tissues of *O. niloticus* exposed to different concentrations of ferric chloride at different period than that control fish tissues. Zinc concentrations in the gill and liver were higher than that in the muscles. Also, there was significant increase in the concentrations of zinc in the gill, liver and muscle tissues of *O. niloticus* exposed to different concentrations of zinc sulfate at different period than that control fish tissues. Results generally showed that metal concentrations were always lowest in the muscles and highest in the gill and liver.

### INTRODUCTION

The freshwater fish has been affected and accumulated the heavy metals in their tissues as reported by Shokr (2007) who indicated that heavy metals accumulated in their tissues due to exposed to heavy metales. Some of the marine environment world-wide has been contaminated by heavy metals and as a result of (Mance, 1987; Langston, 1990; Merian, 1991; Bryan and Langston, 1992) this animals living in contaminated waters showed high metal concentrations in their tissues.

Over a few decades there has been growing interest to determine heavy metal levels in the marine environments and attention was drawn to find out contamination level of public food supplies particularly fish. Therefore, the marine environments are occasionally monitored for heavy metal contamination in water, sediment and animals. It is well known that heavy metals accumulate in tissues of aquatic animals and therefore heavy metals measured in tissues of aquatic animals can reflect the past exposures. Tissue concentrations of heavy metals can also be a reasonable measurement for public health standards and for animals' health point of view. Human population growth and industrial development have been the major causes of coastal contamination around the world during recent years (Caussy *et al.*, 2003). The subsequent accumulation of xenobiotic chemicals in sediments, seawater, or prey organisms has been shown to be directly linked to adverse health effects in both humans and fish (Boening, 2000, Riba *et al.*, 2004 and Ashraf, 2005). Contaminant levels in biotic and abiotic components of an ecosystem can easily be measured using accessible chemical analysis techniques, few reliable diagnostic tools are currently available to fish health professionals to evaluate disease in fish. According to Hrubec *et al.*, (2000), many of the clinical tools used to assess mammalian health are not developed for use in fish. Yet various workers have shown that hematological parameters can provide satisfactory information on the physiological response of fish to environmental stressors for two major reasons, namely, the close association of the circulatory system with the external environment (Houston, 1997, Lohner *et al.*, 2001 and Cazenave *et al.*, 2005). Similarly, histopathological investigations have long been recognized to be reliable biomarkers of stress in fish for several reasons described previously (Teh *et al.*, 1997 and van der Oost *et al.*, 2003). Gill lesions as indicators of exposure to contaminants have previously been used in numerous laboratory and field studies around the world (Dalzell and Macfarlane, 1999; and Oliveira Ribeiro *et al.*, 2002) including Mauritius (Bhagwant and Elahee, 2002). Unlike the enucleated mammalian red blood cells, fish erythrocytes possess very large nuclei, which may occupy about 29% of the cell volume (Wilhelm-Filho *et al.*, 1992 and Shokr, 2011).

The aim of this study is to determine heavy metal (Pb, Zn and Fe) concentrations in the muscles, gills and livers of freshwater fish (*Oreochromis niloticus*) exposed to lead acetate ferric chloride and zinc sulfate. This fish was very important source of food to human being.

## MATERIALS AND METHODS

The *O. niloticus* 80 - 100 gm body weight was obtained from Abbassa farm. The fish were reared in large tanks contained water from their farm. The fish were acclimatized and fed on 30 % protein diet. The fish were divided into four groups, control group and three groups exposed to 5, 10 and 15 mg/l of lead acetate, 10, 20 and 30 mg L<sup>-1</sup> of ferric chloride and 10, 20 and 300 mg L<sup>-1</sup> of zinc sulfate respectively, the experimental fish were exposed to treatment for periods of 15 days, 30 days and 45 days. Each group consists of two-glass aquaria 30 x 60 x 40 cm<sup>3</sup>. Each glass aquarium was contained 10 fish. In the laboratory, they were immediately dissected with clean equipment. Total size and weight of the animals were measured. Since size may be an important factor in heavy metal accumulation of aquatic animals (Canli and Furness, 1993a and b). six fish were obtained from each group after 15, 30 and 45 days. The gill, liver and muscle tissues of the animals were dissected using clean equipment and put in petri dishes. Tissues of the each group were pooled to make 5 subsamples and pooled tissues were transferred to an oven set at 120 °C to

dry. Drying continued until all the wet tissues reached to a constant weight, Dry tissue samples were put into digestion flasks and 4 ml perchloric acid and 8 ml nitric acid (Merck) were added. The digestion flasks were then put on a hot plate set at 120 °C (gradually increased) until all the materials were dissolved. After digestion the digested samples were diluted with distilled water appropriately in the range of standards which were prepared from stock standard solution of the metals (Merck). Metal concentrations in the samples were measured using a Perkin Elmer AS 3100 flame atomic absorption spectrophotometer. Metal concentration in a tissue was presented as  $\mu\text{g}$  metal/gm dry weight.

#### Statistical Analysis:

Data were recorded as means  $\pm$  S. E. (n) significant differences ( $p < 0.05$ ) within each group were tested with Student's two-tailed t – test and one-way ANOVA by SPSS for windows XP2 2002, comparisons between groups were tested by Student's two-tailed t – test unpaired design, ( $p < 0.05$ ).

## RESULTS

Results of Table (1) revealed that the level of lead in gills increased significantly ( $P < 0.05$ ) with each increase in its concentration in water, however concentrations in the liver and muscle showed in significant increase with its increase in water to 5 mg/l compared to the control group and significantly with its increasing level to 10 and 15 mg/l compared to control group and the group that exposed to 5 mg/l of lead acetate with each exposure period.

Table 1: the concentrations of lead ( $\mu\text{g}/\text{gm}$  d. w.) in the tissues of *Oreochromis niloticus* exposed to 5, 10 and 15 mg/l of lead acetate at different period

Organs	Doses /Periods	15 days	30 days	45 days
Gills	Control	3.65 $\pm$ 0.095	3.10 $\pm$ 0.065	3.32 $\pm$ 0.054
	5 mg/l of lead acetate	7.85 $\pm$ 0.042*	8.83 $\pm$ 0.065*	9.25 $\pm$ 0.042*
	10 mg/l of lead acetate	9.24 $\pm$ 0.47**	10.25 $\pm$ 0.089**	12.24 $\pm$ 0.036**
	15 mg/l of lead acetate	12.05 $\pm$ 0.048***	12.25 $\pm$ 0.069***	14.02 $\pm$ 0.95***
Liver	Control	1.33 $\pm$ 0.084	1.24 $\pm$ 0.18	1.81 $\pm$ 0.21
	5 mg/l of lead acetate	4.20 $\pm$ 0.32	5.21 $\pm$ 0.64	6.02 $\pm$ 0.46
	10 mg/l of lead acetate	8.11 $\pm$ 0.27**	9.41 $\pm$ 0.24**	9.22 $\pm$ 0.51**
	15 mg/l of lead acetate	10.02 $\pm$ 0.35***	10.71 $\pm$ 0.24***	12.28 $\pm$ 0.44***
Muscles	Control	1.24 $\pm$ 0.22	1.27 $\pm$ 0.68	1.24 $\pm$ 0.66
	5 mg/l of lead acetate	2.95 $\pm$ 0.58	3.83 $\pm$ 0.14	5.82 $\pm$ 0.57
	10 mg/l of lead acetate	5.35 $\pm$ 0.54**	6.23 $\pm$ 0.26**	8.23 $\pm$ 0.24**
	15 mg/l of lead acetate	7.21 $\pm$ 0.97***	8.69 $\pm$ 0.84***	9.94 $\pm$ 0.64***

In the gills level of iron increased significantly ( $P < 0.05$ ) with each increase in its concentration in water, however concentrations in the liver and muscle showed in significant increase with its increase in water to 10 mg/l compared to the control group and significantly with its increasing level to 20 and 30 mg/l compared to control group and the group that exposed to 10 mg/l of ferric chloride with each exposure period as shown in Table (2).

Zinc concentrations in the gills, liver and muscles of *Oreochromis niloticus* exposed to different concentrations of zinc sulfate at different periods are shown in Table (3). level of Zinc in gills increased significantly ( $P < 0.05$ ) with each increase in its concentration in water, however concentrations in the liver and muscle showed in significant increase with its increase in water to 10 mg/l compared to the control group and significantly with its increasing level to 20 and 30 mg/l compared to control group and the group that exposed to 10 mg/l of zinc sulfate with each exposure period.

Table 2: The concentrations of iron ( $\mu\text{g}/\text{gm d. w.}$ ) in the tissues of *Oreochromis niloticus* exposed to 10, 20 and 30  $\text{mg L}^{-1}$  of ferric chloride at different period

	Periods/organs	10 days	20 days	30 days
Gills	control	61.94±0.58	60.32±0.14	60.61±0.52
	10 mg L-1 of ferric chloride	83.66±0.95***	86.81±0.57***	84.24±0.57***
	20 mg L-1 of ferric chloride	88.23±0.91***	89.25±0.41***	93.35±0.84***
	30 mg L-1 of ferric chloride	90.04±0.52***	93.35±0.42***	94.32±0.81***
Liver	control	48.23±0.51	49.51±0.81	49.21±0.62
	10 mg L-1 of ferric chloride	65.29±0.45	65.36±0.91*	65.56±0.56*
	20 mg L-1 of ferric chloride	80.26±0.41**	82.52±0.95**	83.35±0.47**
	30 mg L-1 of ferric chloride	82.32±0.83***	85.23±0.71***	88.89±0.61***
Muscle	control	50.3±0.51	50.5±0.83	50.3±0.47
	10 mg L-1 of ferric chloride	55.22±0.51	65.24±0.61*	65.89±0.52*
	20 mg L-1 of ferric chloride	63.55±0.71*	71.52±0.61**	75.88±0.91**
	30 mg L-1 of ferric chloride	69.52±0.54**	72.35±0.86***	85.56±0.76***

Table 3: The concentrations of zinc ( $\mu\text{g} / \text{gm d. w.}$ ) in the tissues of *Oreochromis niloticus* exposed to 10, 20 and 30  $\text{mg L}^{-1}$  of zinc sulfate at different period

	Periods/organs	10 days	20 days	30 days
Gills	control	14.3±0.84	14.3±0.51	14.2±0.57
	10 mg L-1 of zinc sulfate	17.11±0.24*	18.21±0.81*	20.25±0.62*
	20 mg L-1 of zinc sulfate	19.85±0.31***	20.98±0.61***	23.87±0.95***
	30 mg L-1 of zinc sulfate	25.21±0.22***	26.57±0.71***	27.98±0.75***
Liver	control	13.2±0.42	13.9±0.71	13.6±0.84
	10 mg L-1 of zinc sulfate	15.88±0.87*	16.52±0.63*	20.3±0.85*
	20 mg L-1 of zinc sulfate	18.22±0.81***	21.35±0.64***	25.6±0.62***
	30 mg L-1 of zinc sulfate	24.21±0.83***	26.11±0.76***	27.55±0.81***
Muscle	control	10.6±0.80	10.2±0.63	10.7±0.41
	10 mg L-1 of zinc sulfate	13.32±0.44*	14.36±0.44*	15.24±0.88*
	20 mg L-1 of zinc sulfate	16.31±0.83***	17.14±0.82***	18.25±0.77***
	30 mg L-1 of zinc sulfate	21.22±0.51***	21.86±0.91***	23.15±0.69***

Significant at  $p < 0.05$ , \*\* highly significant at  $p < 0.01$ , \*\*\* very highly significant at  $p < 0.001$

Results generally showed that metal concentrations were always lowest in the muscles and highest in the gills and liver.

## DISCUSSION

In the present study the fish that exposed to 5, 10 and 15  $\text{mg/l}$  of lead acetate, 10, 20 and 30  $\text{mg L}^{-1}$  of ferric chloride and 10, 20 and 300  $\text{mg L}^{-1}$  of zinc sulfate respectively, showed that metal concentrations were always lowest in the muscles and highest in the gills and liver. This is probably due to their physiological roles in fish metabolism. It has been shown that target tissues of heavy metals are metabolically active ones, like the liver, kidney and gills. Therefore, metal accumulation in these tissues occurs at higher level compared to some other tissues, like the muscles, where metabolic activity is relatively low (Heath, 1987; Langston, 1990; Roesijadi and Robinson, 1994; Canli *et al.*, 1995). It is generally agreed that heavy metal uptake occurs mainly from water, food and sediment. However, effectiveness of metal uptake from these sources may differ in relation to ecological needs and metabolism of animals and also contamination gradients of water, food and sediment as well as some other factors such as salinity, temperature, interacting agents (Heath, 1987; Langston, 1990; Roesijadi and Robinson, 1994). Kilgour (1991) indicated that animals which have close relationship with sediment, show relatively high body concentrations of cadmium, although uptake from water was more important route than uptake from sediment for animals which do not burrow. Canli and Furness (1998) also showed that tissue distribution of metals in the Norway lobster *Nephrops norvegicus* differed

significantly following to an uptake protocol from food and from seawater. Ecological needs of fishes are also one of the most important factors in the accumulation of heavy metal. For example, in the eel *Anguilla anguilla*, whiting *Merlangius merlangius*, flounder *Platichthys flesus* and plaice *Pleuronectes platessa*, levels of Zn, Cu, Pb and Cd in the muscle ranged considerably (Wharfe and Van Der Broek, 1977).

In the present study *O. niloticus* exposed to 5, 10 and 15 mg/l of lead acetate, 10, 20 and 30 mg L<sup>-1</sup> of ferric chloride and 10, 20 and 300 mg L<sup>-1</sup> of zinc sulfate respectively, for a period of 20 days, 30 days and 45 days this different from ecological needs of the fishes and play an outstanding role as all the fish species regardless of their ecological needs showed high metal concentrations. This may indicate that the exposure of *O. niloticus* to 5, 10 and 15 mg/l of lead acetate, 10, 20 and 30 mg L<sup>-1</sup> of ferric chloride and 10, 20 and 300 mg L<sup>-1</sup> of zinc sulfate respectively, for a period of 20 days, 30 days and 45 days causes increase precipitation of heavy metals as lead, iron and zinc in the tissues of this fish as gills, liver and muscles and increase the precipitation in the gills and liver than that in the muscle. Data from the literature showed that metal concentrations in the tissues of fish varied widely depending on where the animals were caught and exposure to heavy metals toxicity. Doganoc (1995) found that fishes from Slovenia contained (ppm w.w.) <0.05-0.34 Pb, <0.003-0.05 Cd, in the muscle between the period of 1982-1993. Serra et al. (1993) showed that in tissues of *Liza ramada* and *Leuciscus cephalus* fished from the northern Adriatic, zinc levels were higher in the chubb than the grey mullet. They also found that the concentrations of iron and copper in the red muscle of the grey mullets were remarkably high and they indicated that this was probably due to swimming activity of this species. Remarkably high lead, iron and zinc concentrations in the liver of the *O. niloticus* in the present study support the high lead, iron and zinc concentrations of this species. Medina et al. (1986) measured the concentrations of heavy metals in the muscle of *Mullus barbatus* and *Mullus surmelatus*, collected from Spanish coasts. They found that Cd, Pb and Cr levels ranged between 0.019-0.187, 0.03-0.803 and 0.023-0.638 µg/g d.w. These values are lower when compared to the present results. Iron and zinc concentrations also may vary between different fish. For example, iron and zinc concentrations in the muscles of various marine fishes from Bangladesh ranged between 8.50-22.2 ppm w.w. for iron and 7.40-22.5 ppm w.w. for zinc (Quazi et al. 1995). The present study also showed that iron and zinc concentrations also varied among the groups of fishes that exposed to acute toxicity of ferric and zinc. This is perhaps due to the fishes that exposed to acute toxicity of ferric and zinc. Romeo (1987) also found relatively low concentrations of Cu, Zn, Cd and Pb in the muscle of *Mugil cephalus* (mullet roe) from the northern coast of Mauritania in the Atlantic Ocean as 2.3, 142, <0.1, <0.5 ppm d.w. respectively. However, this is not surprising because oceans are less contaminated marine environments comparing to seas that generally face human impact more than oceans. This study also emphasized that some metal levels were higher than the acceptable values for human consumption set by various health organizations. For example, zinc, iron and Pb data given in the present study are considerably high compared to daily tolerable cadmium and lead intake from food according to WHO/FAO committee's proposal (Merian, 1991). These are approximately 1 µg Cd/kg body weight and 7 µg Pb/kg body weight. This study showed that heavy metal concentrations in the tissues of fishes, regardless of their exposure levels and periods, were considerably high at all groups compared to metal concentrations in the literature given above from other waters. So, there is a high risk

for consuming this fish to human being, and treatment of water is highly recommended before using it in aquaculture.

#### ACKNOWLEDGEMENT

The researcher thanks the Central Lab of Aquaculture Research, Abbasa for helping in performing this study.

#### REFERENCE

- Ashraf, (2005). Accumulation of heavy metals in kidney and heart tissues of Epinephelus microdon fish from the Arabian Gulf. *Environ. Monit. Assess.*, 101: 311–316.
- Bhagwant, and Elahee, K.B. (2002). Pathologic gill lesions in two edible fish species, *Mulloidichthys flavolineatus* and *Mugil cephalus* from the Bay of Poudre d'Or, Mauritius. *WIO J. Mar. Sci.*, 1: 35–42.
- Boening, Ecological effects, transport, and fate of mercury: A general review. *Chemosphere*, 40: 1335–1351.
- Bryan and Langston WJ. (1992). Bioavailability, accumulation and effects of heavy metals in sediments with special reference to United Kingdom estuaries: a review. *Environ Pollut.*, 76: 89-131.
- Canli and Furness RW. (1993a). Heavy metals in tissues of the Norway lobster *Nephrops norvegicus*: Effects of sex, size and season, *Chem. Ecol.*, 8: 19-32.
- Canli and Furness RW. (1995). Mercury and cadmium uptake from seawater and from food by the Norway lobster *Nephrops norvegicus*. *Environ Toxicol. Chem.*, 14: 819-828.
- Canli and Furness RW. (1993b). Toxicity of heavy metals dissolved in sea water and influences of sex and size on metal accumulation and tissue distribution in the Norway lobster *Nephrops norvegicus*. *Mar Environ Res.*, 36: 217-236.
- Canli Ay Ö and Kalay M. (1998). Levels of heavy metals (Cd, Pb, Cu, Cr and Ni) in tissues of *Cyprinus carpio*, *Barbus capito* and *Chondrostoma regium* from the Seyhan River, Turkey. *Turkish J. Zoology.*, 22: 149-157.
- Caussy, Gochfeld, M., Gurzau, E., Neagu, C. and Ruede, H. (2003). Lessons from case studies of metals: investigating exposure, bioavailability and risk. *Ecotoxicol. Environ. Safety*, 56: 45–51.
- Cazenave, Wunderlin, D. A., Hued, A. C. and de los Angeles-Bistoni, M. (2005). Haematological parameters in a neotropical fish, *Corydoras paleatus* (Jenyns, 1842) (Pisces, Callichthyidae), captured from pristine and polluted water. *Hydrobiologia*, 537: 25–33.
- Dalzell, and Macfarlane, N. A. A. (1999). The toxicity of iron to brown trout and effects on the gills: A comparison of two grades of iron sulphate. *J. Fish Biol.* 55:301–315.
- Doganoc (1995). Heavy metals concentrations in freshwater and marine fishes in period 1982-1993. *Zbornik Biotehniške Fakultete Univerze v Ljubljani (Slovenia). Kmetijstvo (Zootehnika)* 66: 89-97.
- Heath (1987). *Water pollution and fish physiology*. CRC press, Florida, 245 pp.
- Hematological and population level assessment. *Ecotoxicol. Environ. Safety* (2001). 50: 203–216.
- Houston, (1997). Are the classical haematological variables acceptable indicators of fish health? *Trans. Am. Fish. Soc.*, 126:879–894.

- Hrubec, Cardinale, J. L. and Smith, S. A. (2000). Haematology and plasma chemistry reference intervals for cultured tilapia (*Oreochromis hybrid*). *Vet. Clin. Pathol.*, 29: 7–12.
- Kilgour (1991). Cadmium uptake from cadmium-spiked sediments by four freshwater invertebrates. *Bull Environ Contam Toxicol*, 47:70-75.
- Langston (1990). Toxic effects of metals and the incidence of marine ecosystems. In: Furness RW, Rainbow PS, editors. *Heavy Metals in the Marine Environment*. CRC Press, New York, 256 pp.
- Lohner, Reash, R. J., Willet, V.E. and Rose, L.A., Assessment of tolerant sunfish populations (*Lepomis sp.*) inhabiting selenium-laden coal ash effluents. 1.
- Mance (1987). *Pollution threat of heavy metals in aquatic environment*. Elsevier, London, 363 pp.
- Medina, Hernandez F, Pastor A, Beyerfull JB and Barbera JC. (1986). Determination of mercury, cadmium, chromium and lead in marine organisms by flameless atomic absorption spectrophotometry. *Mar Pollut Bull*, 17:41-44.
- Merian (1991). *Metals and Their Compounds in the Environments. Occurrence, Analysis and Biological Relevance*. ISBN 0-89573 - 562 - 8 (VCH New York).
- Oliveira Ribeiro, Belger, L., Pelletier, E. and Rouleau, C. (2002). Histopathological evidence of inorganic mercury and methyl mercury toxicity in the Arctic charr (*Salvelinus alpinus*). *Environ. Res.*, 90: 217–225.
- Quazi Banu CP, Rahman MM and Sayeed S. (1995). Mineral content of fresh water and marine fish species. *Proceedings of the third Asia-Pacific food analysis Network Conference on Food Analysis*. P. 2, (IRRI call no. TX511 C65).
- Riba, Conradi, M., Forja, J.M. and DelValls, T.A., (2004). Sediment quality in the Guadalquivir estuary: Lethal effects associated with the Aznalcollar mining spill. *Mar. Pollut. Bull.*, 48:144–152.
- Roesijadi and Robinson WE. (1994). Metal regulation in aquatic animals: Mechanism of uptake, accumulation and release. In: Malins DC, Ostrander GK, editors. *Aquatic Toxicology (Molecular, Biochemical and Cellular Perspectives)*, Lewis Publishers, London, 539 pp.
- Romeo Trace metals in fish roe from the Mauritania coast. *Mar Pollut Bull*(1987).18:507-508.680
- Serra, Carpena E, Torresani G, Andreucci A and Grandini S. (1993). Concentrations of Zn, Cu, Fe, and Cd in *Liza ramada* and *Leuciscus cephalus*. *Archivio Veterinario Italiano*, 44:166-174.
- Shokr, (2011). Hematological and Biochemical Parameters of Nile Tilapia; *Oreochromis niloticus* under Pollution of Iron. *Egypt. J. Aquat. Biol. & Fish.*, 15(3): 33- 47.
- Shokr: Accumulation of Aluminum in Some Organs of *Tilapia zillii* G. Exposed to Aluminum Chloride and Physiological Changes in Serum and Liver. *Egypt. J. Appl. Sci.*, (2007) .22 (9): 1 – 10
- Teh, Adams, S.M. and Hinton, D.E. (1997). Histopathologic biomarkers in feral freshwater fish populations exposed to different types of contaminant stress. *Aquat. Toxicol.*, 37:51–70.
- Van der Oost, Beyer, J and Vermeulen, N.P.E. (2003). Fish bioaccumulation and biomarkers in environmental risk assessment: A review. *Environ. Toxicol. Pharmacol.* 13:57–149.
- Wharfe Van don and Broek WLF. (1977). Heavy metals in macroinvertebrates and fish from the lower Medway Estuary, Kent. *Mar Pollut Bull.*, 8: 31-35.

Wilhelm-Filho, Eble, G.J., Kassner, G., Caprario, F.X. and Dafre, A. L., (1992).  
Blood cells from trout. J. Fish Biol., 59: 1098–1103.

## ARABIC SUMMERY

### المعادن الثقيلة (الرصاص و الحديد و الزنك) فى انسجة الاسماك نتيجة لتلوث المياه

السيد احمد محمد شكر

قسم الفسيولوجى- كلية الطب- جامعة حائل

اوضحت النتائج ان تعرض اسماك البلطى النيلية الى تركيزات مختلفة و اوقات مختلفة من اسيتات الرصاص الى ارتفاع فى مستوى الرصاص فى انسجة البلطى النيلية المختلفة من الخياشيم و الكبد و العضلات. وجد ارتفاع فى مستوى تركيز الرصاص فى الخياشيم و الكبد اكثر من العضلات تحت تاثير الجرعات المختلفة من اسيتات الرصاص.

اظهرت النتائج ان تعرض اسماك البلطى النيلية الى تركيزات مختلفة و اوقات مختلفة من كلوريد الحديد الى ارتفاع فى مستوى الحديد فى انسجة البلطى النيلية المختلفة من الخياشيم و الكبد و العضلات. وجد ارتفاع فى مستوى تركيز الحديد فى الخياشيم و الكبد اكثر من العضلات تحت تاثير الجرعات المختلفة من كلوريد الحديد.

اوضحت النتائج ان تعرض اسماك البلطى النيلية الى تركيزات مختلفة و اوقات مختلفة من كبريتات الزنك الى ارتفاع فى مستوى الزنك فى انسجة البلطى النيلية المختلفة من الخياشيم و الكبد و العضلات. وجد ارتفاع فى مستوى تركيز الرصاص فى الخياشيم و الكبد اكثر من العضلات تحت تاثير الجرعات المختلفة من كبريتات الزنك.